

Transport in Humans

Matthew Williams • Biology • May 9, 2026

Transport in Humans

As organisms grow larger, diffusion alone cannot deliver substances fast enough — the distance from the surface to the innermost cells becomes too great. Multicellular organisms therefore develop specialised transport systems to move materials quickly over long distances.

Why Transport Systems Are Needed


A small organism has a high surface area to volume ratio, so diffusion supplies every cell adequately. In a large organism the ratio is much lower, and cells deep inside would starve of oxygen or accumulate waste. Transport systems solve this by:

- delivering oxygen, glucose, and hormones to every tissue
- removing carbon dioxide, urea, and other metabolic wastes
- distributing heat evenly around the body

The Human Circulatory System

Humans have a **closed, double circulatory system**:

- **closed** — blood stays within blood vessels at all times
- **double** — blood passes through the heart twice per complete circuit (once for the lungs, once for the body)



Double circulation in humans

Pulmonary circulation: right ventricle 'lungs 'left atrium. Blood picks up oxygen and loses CO,

Systemic circulation: left ventricle 'body 'right atrium. Blood delivers oxygen and nutrients; returns with CO, and waste.

The Heart

The heart is a muscular pump with four chambers: two atria (receive blood) and two ventricles (pump blood out). Valves prevent backflow.

- **Right side** receives deoxygenated blood from the body and pumps it to the lungs
- **Left side** receives oxygenated blood from the lungs and pumps it to the body
- The left ventricle has thicker walls than the right because it pumps blood the greater distance around the entire body

Labelled diagram of the human heart showing atria, ventricles, valves, aorta, vena cava, and pulmonary vessels

Blood Vessels

Vessel	Wall structure	Lumen	Valves	Direction	Function
Artery	thick, muscular, elastic	narrow	no	away from heart	carries blood at high pressure
Vein	thin, less muscular	wide	yes	toward heart	returns blood at low pressure

Capillary	one cell thick	very narrow	no	connects arteries to veins	site of exchange between blood and tissues
-----------	----------------	-------------	----	----------------------------	--

Capillaries are so thin that red blood cells must squeeze through in single file. Their walls allow plasma to leak out, forming tissue fluid that bathes cells.

Blood

Blood is a liquid connective tissue. It has four main components:

Plasma

Plasma is the pale yellow liquid portion of blood (about 55%). It transports:

- dissolved glucose, amino acids, fatty acids
- hormones
- CO₂ (partly as hydrogencarbonate ions)
- urea
- heat

Red Blood Cells (Erythrocytes)

Red blood cells carry oxygen using **haemoglobin**, an iron-containing protein that combines reversibly with oxygen:

haemoglobin + oxygen → oxyhaemoglobin

In the lungs (high O₂): oxyhaemoglobin forms. In the tissues (low O₂): oxygen is released.

Adaptations of red blood cells:

- **biconcave disc shape** — large surface area relative to volume; short diffusion distance to centre
- **no nucleus** — maximum space for haemoglobin
- **flexible membrane** — squeezes through narrow capillaries
- produced in vast numbers — high oxygen-carrying capacity

White Blood Cells (Leucocytes)

White blood cells defend the body. Two main types:

Phagocytes — engulf and destroy pathogens by phagocytosis (the phagocyte surrounds the pathogen, encloses it in a vacuole, and enzymes from lysosomes digest it).

Lymphocytes — produce antibodies. B-lymphocytes differentiate into:

- **plasma cells** — secrete antibodies specific to the antigen
- **memory cells** — remain in the blood long-term; allow a much faster response to the same pathogen on re-exposure

T-lymphocytes coordinate the immune response and attack infected or abnormal cells.

Platelets

Platelets are cell fragments that trigger blood clotting when a vessel is damaged:

damaged tissue 'platelets release thromboplastin 'prothrombin 'thrombin 'fibrinogen 'fibrin mesh 'clot forms

Clotting prevents excessive blood loss and blocks entry of pathogens.

Immunity and Vaccination

Antigens are foreign molecules (usually on pathogen surfaces) that trigger an immune response.

Antibodies are proteins produced by B-lymphocytes that are specific to one antigen. They may neutralise toxins, cause pathogens to clump (agglutination), or mark them for destruction.

Type of immunity	How acquired	Duration
Active natural	body produces own antibodies after infection	long-lasting
Active artificial	body produces own antibodies after vaccination	long-lasting
Passive natural	antibodies passed from mother to baby via placenta or breast milk	temporary
Passive artificial	ready-made antibodies injected	temporary

Vaccination works by introducing weakened or dead pathogens, or their antigens, so the immune system produces antibodies and memory cells without the person suffering the disease. If the pathogen is later encountered, the memory cells allow a rapid secondary immune response.

 **Exam Tip**

Active immunity lasts because memory cells are produced. Passive immunity is immediate but short-lived because no memory cells are made — the antibodies eventually break down.

PRACTICE — TRANSPORT IN HUMANS**Double circulation**

A circulatory system in which blood passes through the heart twice per circuit: once to the lungs, once to the body.

Artery

A thick-walled blood vessel that carries blood away from the heart at high pressure.

Vein

A thin-walled blood vessel with valves that returns blood to the heart at low pressure.

Capillary

A vessel one cell thick; site of exchange between blood and body tissues.

Haemoglobin

The iron-containing protein in red blood cells that reversibly binds oxygen.

Phagocytosis

The process by which white blood cells engulf and destroy pathogens.

Antibody

A protein produced by B-lymphocytes that is specific to one antigen and helps destroy pathogens.

Active immunity

Long-lasting immunity in which the body produces its own antibodies and memory cells.

Passive immunity

Short-lived immunity from receiving ready-made antibodies; no memory cells produced.