# Averages, Spread & Cumulative Frequency

Matthew Williams • Math • May 6, 2026

Averages describe the centre of a dataset, while spread describes how far the values are from each other. Two classes can have the same mean score but very different consistency, so both centre and spread are needed to understand the data properly.

CSEC questions often move from calculation to interpretation: find the mean, median, range, quartiles, or cumulative frequency, then say what the result tells you. Do not treat these as isolated formulas. After calculating, write a short sentence explaining what the number means in the context of the question.

A **measure of central tendency** represents the "typical" or "average" value in a dataset.

## Mean (Average)

The mean uses every value, so it is affected by very large or very small outliers. It is useful when the data is fairly balanced.

$$\text{Mean} = \frac{\text{Sum of all values}}{\text{Number of values}}$$

**For raw data:**

📋 **Example**

**Scores: 5, 7, 8, 9, 5**

$$\text{Mean} = \frac{5+7+8+9+5}{5} = \frac{34}{5} = 6.8$$

**For ungrouped frequency data:**

$$\text{Mean} = \frac{\sum(value \times frequency)}{\sum frequency}$$

📋 **Example**

| Score | Frequency |
|---|---|
| 5 | 2 |
| 7 | 1 |
| 8 | 1 |
| 9 | 1 |

$$\text{Mean} = \frac{(5 \times 2) + (7 \times 1) + (8 \times 1) + (9 \times 1)}{2 + 1 + 1 + 1} = \frac{10 + 7 + 8 + 9}{5} = \frac{34}{5} = 6.8$$

**For grouped data:**

Use class midpoints:

$$\text{Mean} = \frac{\sum(\text{midpoint} \times \text{frequency})}{\sum \text{frequency}}$$

📋 **Example**

| Class | Frequency | Midpoint |
|---|---|---|
| 10-19 | 3 | 14.5 |
| 20-29 | 5 | 24.5 |
| 30-39 | 2 | 34.5 |

$$\text{Mean} = \frac{(14.5 \times 3) + (24.5 \times 5) + (34.5 \times 2)}{3 + 5 + 2}$$

$$= \frac{43.5 + 122.5 + 69}{10} = \frac{235}{10} = 23.5$$

## Median

The median is the middle value after ordering the data. It is useful when outliers would distort the mean.

The **median** is the middle value when data is arranged in order.

**For raw data:**

- 1. Arrange values from smallest to largest
- 2. If odd number of values: median is the middle one

- 3. If even number of values: median is the average of the two middle ones

> 📋 **Example**
> **Scores: 5, 5, 7, 8, 9 (5 values, odd)**
> Median = 7 (the 3rd value)
> **Scores: 5, 5, 7, 8, 9, 10 (6 values, even)**
> Median = (7 + 8) ÷ 2 = 7.5

**For grouped data:**

Use the cumulative frequency table and interpolation:

$$\text{Median} = L + \frac{\frac{n}{2} - CF}{f} \times w$$

Where:

- $L$ = lower boundary of median class
- $n$ = total frequency
- $CF$ = cumulative frequency before median class
- $f$ = frequency of median class
- $w$ = class width

> 📋 **Example**
> | Class | Frequency | Cumulative |
> |---|---|---|
> | 10-19 | 3 | 3 |
> | 20-29 | 5 | 8 |
> | 30-39 | 2 | 10 |
>
> Total = 10, so median position = 10÷2 = 5
> Median class is 20-29 (cumulative frequency reaches 5 here)
>
> $$\text{Median} = 19.5 + \frac{5-3}{5} \times 10 = 19.5 + \frac{2}{5} \times 10 = 19.5 + 4 = 23.5$$

## Mode

The mode identifies the most common value. It is especially useful for categorical data, where mean and median may not make sense.

The **mode** is the value that appears most often.

> 📋 **Example**
>
> **Scores: 5, 5, 5, 7, 8, 9, 9**
>
> Mode = 5 (appears 3 times)
>
> **Data: 2, 5, 5, 7, 7, 9**
>
> Two modes: 5 and 7 (both appear twice) = **bimodal**
>
> **Data: 2, 5, 7, 9**
>
> No mode (all appear once) = **no mode**

## Choosing Mean, Median, or Mode

Choosing the average is a reasoning skill. The best measure depends on the shape of the data and what the question is trying to describe.

**Use MEAN when:**

- Data is roughly symmetric
- No extreme outliers
- You want to use all values
- Example: class average on a test

**Use MEDIAN when:**

- Data has outliers or is skewed
- You want the "typical" middle value
- Example: house prices (skewed by luxury homes)

**Use MODE when:**

- Categorical data (colors, preferences)
- Discrete data with clear peaks
- Example: favorite color, most common shoe size

📝 **Exam Tip**

**Example: Which average?**

House prices: 120,000, 125,000, 130,000, 140,000, 2,000,000

- Mean = (120+125+130+140+2000)÷5 = **502,300** (way too high!)
- Median = **130,000** (better,the luxury home is an outlier)
- Mode = no mode

**Best answer: Median**, because the data has an outlier.

## Measures of Spread (Dispersion)

**Spread** measures how far apart the data values are from each other.

### Range

Range gives a quick sense of spread, but it only uses the smallest and largest values. One unusual value can make the range misleading.

$$\text{Range} = \text{Maximum value} - \text{Minimum value}$$

📋 **Example**

**Scores: 5, 7, 8, 9, 5**

Range = 9 - 5 = 4

**Problem:** Only uses the extreme values. Doesn't show middle spread.

### Quartiles and Interquartile Range

Quartiles split ordered data into four parts. The interquartile range focuses on the middle half of the data, so it is less affected by extremes.

**Quartiles** divide the data into 4 equal parts.

- $Q_1$ (1st quartile) = 25th percentile
- $Q_2$ (2nd quartile) = 50th percentile = **median**
- $Q_3$ (3rd quartile) = 75th percentile

**Interquartile Range (IQR):**

$$\text{IQR} = Q_3 - Q_1$$

This shows the spread of the middle 50% of data.

📋 **Example**

**Test scores: 5, 6, 7, 7, 8, 8, 8, 9, 9, 10 (10 values)**

Arrange in order: 5, 6, 7, 7, 8, 8, 8, 9, 9, 10

**Q** position = (10+1) ÷ 4 = 2.75 'between 2nd and 3rd values = 6 + 0.75(7-6) = 6.75

**Q** position = (10+1) ÷ 2 = 5.5 'between 5th and 6th values = 8

**Q** position = 3(10+1) ÷ 4 = 8.25 'between 8th and 9th values = 9 + 0.25(9-9) = 9

$$\text{IQR} = Q_3 - Q_1 = 9 - 6.75 = 2.25$$

## Semi-Interquartile Range

The semi-interquartile range is half of the IQR. It gives a compact measure of spread around the middle of the dataset.

$$\text{Semi-IQR} = \frac{\text{IQR}}{2} = \frac{Q_3 - Q_1}{2}$$

📋 **Example**

Using the example above:

$$\text{Semi-IQR} = \frac{2.25}{2} = 1.125$$

🧠 **Remember**

- **Range**: Uses only extremes, sensitive to outliers
- **IQR**: Shows spread of middle 50%, ignores outliers
- **Semi-IQR**: Half of IQR, useful for comparison

# Cumulative Frequency and Ogives

## Cumulative Frequency Table

Cumulative frequency is a running total. It answers questions like "how many values are less than or equal to this point?"

**Cumulative frequency** = total count up to and including that class.

> 📋 **Example**
> | Class | Frequency | Cumulative Frequency |
> |---|---|---|
> | 10-19 | 3 | 3 |
> | 20-29 | 5 | 3+5 = 8 |
> | 30-39 | 7 | 8+7 = 15 |
> | 40-49 | 4 | 15+4 = 19 |
> | 50-59 | 1 | 19+1 = 20 |

## Cumulative Frequency Curve (Ogive)

An ogive turns cumulative totals into a graph. It is useful for estimating medians, quartiles, and percentiles from grouped data.

An **ogive** is an S-shaped curve showing cumulative frequency.

**How to draw:**

- 1. Use class boundaries on x-axis (not class limits)
- 2. Use cumulative frequency on y-axis
- 3. Plot point at upper boundary of each class
- 4. Connect points with a smooth curve

📋 **Example**

Using the table above:

| Upper Boundary | Cumulative Frequency |
|---|---|
| 19.5 | 3 |
| 29.5 | 8 |
| 39.5 | 15 |
| 49.5 | 19 |
| 59.5 | 20 |

## Reading from an Ogive

To read from an ogive, move horizontally from the cumulative frequency value to the curve, then down to the data value. This is an estimate, so use the graph carefully.

You can read:

- **Quartiles**: $Q_1$ at 25% of total, $Q_2$ at 50%, $Q_3$ at 75%
- **Percentiles**: any value's percentage position
- **Median**: where cumulative frequency = n/2
- **Frequencies above/below** a given value

📋 **Example**

**From the ogive above (n=20):**

**$Q_1$** (25% of 20 = 5): Read across from cumulative frequency 5 to curve, then down to x-axis = 22

**Median** (50% of 20 = 10): Read from cumulative 10 = 32

**$Q_3$** (75% of 20 = 15): Read from cumulative 15 = 39.5